

A service of the National Library of Medicine
and the National Institutes of Health

www.pubmed.gov

My NCBI
[Sign In] [Reg]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

E

Search PubMed for warts podophyllin benzoin

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-UtilitiesPubMed Services
Journals Database
MeSH Database
Single Citation
Matcher
Batch Citation
Matcher
Clinical Queries
Special Queries
LinkOut
My NCBIRelated
Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central**Note:** Performing your original search, *warts podophyllin benzoin*, in PubMed will retrieve **10** citations.

Display Abstract Show 20 Sort by Send to

All: 1 Review: 0

1: Br J Vener Dis. 1981 Jun;57(3):208-9. Related Articles, Links

Podophyllin 10% and 25% in the treatment of ano-genital warts. A comparative double-blind study.

Simmons PD.

One hundred and forty male patients with ano-genital warts were randomly allocated to a double-blind study of 10% and 25% podophyllin in tincture of benzoin compound. One hundred and nine patients attended for a three-month period of surveillance. Only 24 (22%) patients were free of warts after having podophyllin treatment alone, 12 each after treatment with 10% and 25% podophyllin. There was no significant difference in the number of applications needed with each treatment. Neither hypersensitivity nor chemical ulceration occurred.

Publication Types:

- Clinical Trial
- Randomized Controlled Trial

PMID: 7016248 [PubMed - indexed for MEDLINE]

A service of the National Library of Medicine
and the National Institutes of HealthMy NCBI
[Sign In] [Reg]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

E

Search PubMed for warts cantharidin podophyllin Go Clear

Limits

Preview/Index

History

Clipboard

Details

About Entrez
NCBI Toolbar

Text Version

Entrez PubMed
Overview
Help | FAQ
Tutorials
New/Noteworthy
E-UtilitiesPubMed Services
Journals Database
MeSH Database
Single Citation
Matcher
Batch Citation
Matcher
Clinical Queries
Special Queries
LinkOut
My NCBIRelated
Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central**Note:** Performing your original search, **warts cantharidin podophyllin**, in PubMed will retrieve **3 citations**.

Display Abstract Show 20 Sort by Send to

All: 1 Review: 0

1: Pediatr Dermatol. 1984 Jul;2(1):71-3.

Related Articles, Links

Treatment of plantar warts in children with a salicylic acid-podophyllin-cantharidin product.**Coskey RJ.**

One hundred and twenty-one children with plantar warts were treated with a topical preparation containing salicylic acid, 30%; podophyllin, 5%; and cantharidin, 1%. One hundred patients or their families were reached 6 months to 1 year later. This treatment was effective in 81 cases.

PMID: 6504780 [PubMed - indexed for MEDLINE]

Display Abstract Show 20 Sort by Send to

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

May 9 2006 14:13:00